



(405) 235-0040

www.oklahoallergy.com

750 N.E. 13th
3rd Floor

(2 Blocks East of Lincoln Blvd.)
Oklahoma City, Oklahoma 73104-5051

NORTHWEST OFFICE:

Meridian Medical Tower
13321 N. Meridian, Suite 100
Oklahoma City, Oklahoma

NORMAN OFFICE:

Physicians and Surgeons Bldg.
950 North Porter, Suite 102
Norman, Oklahoma

EDMOND OFFICE:

Sycamore Square
120 North Bryant, Suite A4
Edmond, Oklahoma

SPECIALIZING IN THE EVALUATION
AND MANAGEMENT OF
ALLERGIES AND ASTHMA
IN ADULTS AND CHILDREN

John R. Bozalis, MD*

Warren V. Filley, MD*

James R. Clafflin, MD*

Patricia I. Overhulser, MD*

Dean A. Atkinson, MD*

Richard T. Hatch, MD*

Shahan A. Stutes, MD*

Gregory M. Metz, MD*

Karen L. Gregory, CNS

SENIOR CONSULTANTS

Robert S. Ellis, MD*

Charles D. Haunschild, MD*

James H. Wells, MD*

* Diplomate American Board
Allergy and Immunology

Garyl Geist

Chief Operating Officer



URTICARIA (HIVES)/ ANGIOEDEMA (TISSUE SWELLING)

Most “hives” are caused by histamine and other mediators “released” in the upper layers of the skin.

Histamine can be released from special cells, mast cells and basophils, which are found in everyone’s skin.

Patients with hives release “too much” histamine when it is not needed and are classified as:

- “**Acute hivers**” only when they take in an identifiable “trigger”, i.e., drug, food, etc. or if hives last less than 6 weeks.
- “**Chronic hivers**” daily or almost daily without an identifiable “trigger” and lasting for greater than six weeks.

In “chronic” urticaria or persistent hives, a “**trigger**” is rarely found. A “complete” history is needed to look for identifiable “triggers”.

The etiology for chronic urticaria such as medication, heat, pressure or infection is found in less than 5% of patients.

For most cases of chronic urticaria the etiology will remain idiopathic (cause unknown) although an autoimmune etiology can sometimes be found.

- **Angioedema** (tissue swelling), occurs in 90% of patients with chronic urticaria

For chronic urticaria laboratory evaluation can be performed: Thyroid auto-antibody profile, CU Index, complete metabolic profile, complete blood count and autoimmunity tests.

Fortunately, most chronic hives go away with or without treatment:

- 50% of patients hives go away in 3-12 months
- 20% of patients hives go away in 12-36 months
- 20% of patients hives go away in 36-60 months
- 1.5% of patients can have hives for up to 25 years

(60% of “chronic” hivers have recurrences of the hives)

Management is directed toward keeping patients “comfortable” with or without some hives being present. This can be achieved with “appropriate” antihistamine therapy, and occasionally with “other” medications. Prednisone (steroid) should be avoided if at all possible.

Remember the potency of:

- Benadryl 50mg. is considered.....a strength of 1.
- Clarinex, Claritin, Allegra.....a strength of 2-3.
- Atarax (hydroxyzine 25mg.), Zyrtec, Xyzal....a strength of 88.
- Doxepin (Sinequan) 25mg.....a strength of 779.